

Aims

- To investigate changes in VOCs (volatile organic compounds) due to iron supplementation between Day 1 and Day 28 of the study. This will assess VOC changes due to iron supplementation.
- To investigate changes in VOCs following lactulose ingestion. This will assess VOC changes along the gastrointestinal tract.

1. Background and Objectives

Iron deficiency anaemia (IDA) affects approximately >1.2 billion people worldwide^{1,2,3}. In the UK, it can be the reason for up to 13% of referrals to gastroenterologists⁴. Furthermore, The World Health Organisation recognizes IDA as one of the most expensive diseases due to its negative impact on productivity.

IDA can be treated with both oral supplements or IV infusions which are both effective at restoring iron levels in patients. Unabsorbed iron can have unintended side-effects such as enriching

intestinal bacteria that result in bloating due to production of gases. These gases can diffuse into the lungs via the blood and are then detectable on exhaled breath.

Hydrogen and methane are two gases that have been associated with IDA following consumption of the fermentable carbohydrate, lactulose⁵. This research aims to extend this knowledge by exploring whether other gases, volatile organic compounds (VOC), are associated with oral iron supplementation using the lactulose test.

2. Methods

This project was based on VOC changes caused by 28 days of iron supplementation in healthy volunteers.

Owlstone Medical and The Functional Gut Clinic (FGC) were interested in identifying novel breath biomarkers that change in response to oral iron supplementation, and whether production of these biomarkers are related to intestinal geography.

Samples were collected at multiple time points before and after the iron supplementation process, thus each subject served as their own control. Breath samples were analysed at Owlstone Medical Inc. using SIFT-MS technology. Targeted analyses were performed, and compounds deemed statistically significant if they were more than two standard deviations from the lab ambient


IRON SUPPLEMENT:

LACTULOSE CHALLENGE:

BREATH COLLECT:

TIME (MIN):

DAY:



Day	Description	Sampling Time Point (n)			
		T=0 Baseline Before Lactulose Challenge	45 min	90 min	180 min
1	Post-iron suppl.	N=25	N=25	N=25	N=25
28	Post-iron suppl.	N=25	N=25	N=25	N=25

Figure 1: Experimental Design: This was a single-centre, longitudinal study with a population of healthy volunteers monitored before and after exposure to iron supplementation [ClinicalTrials.gov identifier (NCT number): NCT04705662]. 25 adult healthy volunteers were recruited for breath sampling for breath collection using polyvinylidene difluoride (PDVF) breath bags. The site of volunteer induction and sample collection was The Functional Gut Clinic, Manchester. Each volunteer underwent sampling on day 1 before and after administration of lactulose to measure baseline of fermentation levels. After the day 1 visit, volunteers took iron supplements daily and kept a record of any gastrointestinal (GI) tract symptoms experienced. Each volunteer underwent sampling on day 28 ± 2d or sooner if GI symptoms were severe (follow-up clinic visit) before and after administration of lactulose to measure follow-up levels of fermentation.

3. Results

From the 25 healthy volunteers that participated in this study, 2 were excluded due to incomplete samples. Ambient (blank) samples were collected, but not

for all patients/visits/timepoints. Data was symmetrically distributed therefore further mathematical transformation was unnecessary.




Figure 2: PCA analysis showing breath samples are distinct from both lab and room ambient samples using targeted compounds. Pentanoic, butanoic, propanoic, acetic 3-methylbutanoic and hexanoic acids, ethanol, hydrogen sulphide, methane, indole, isoprene, cresols, 2,3-butanedione, trimethylamine, acetone, limonene and phenol were selected in targeted analysis. Lab ambient and room ambient samples show divergent composition, with room ambient resembling the composition of breath samples more than lab samples.

Compound	Adjusted difference	p-value
3-methylbutanoic acid	0.675	0.017
butanoic acid	15.486	0.047
propanoic acid	4.707	0.026
2,3-butanedione	9.793	0.045
limonene	1.307	0.007
hydrogen sulfide	-22.667	0.026
cresol	0.194	0.005

Table 1: Table shows compounds that significantly change after iron supplementation. Linear Mixed models were fitted to evaluate the evolution overtime of the compounds' intensities. These models allow evaluation of both the effect of iron supplementation and that of lactulose challenge, also accounting for the observations' dependence due to repeated measurements from the same HV. For iron supplementation, the model shows that baseline, i.e. time pt. 0, levels of several compounds are significantly different between Day 1 and Day 28.




Figure 3: Graphs showing compounds with significant changes after iron supplementation and/or lactulose challenge. For both (A) propanoic and (B) acetic acids, the trends over time are significant and also different by day of the challenge (either 1 or 28). (C) Hydrogen sulphide, however, significantly decreases after 28 days of iron supplementation. This change is not affected by lactulose ingestion. *p value <0.05.

4. Conclusions

Some short chain fatty acids (SCFAs), butanoic, propanoic and acetic acids, increased after 28 days of iron supplementation following lactulose ingestion: Increases in SCFA has been linked to increased gut health⁵. They have been shown to maintain colonocyte development, promote metabolic health and speculated to play a key role in neuro-immunoendocrine regulation^{6,7}. The significant increase in these SCFAs indicate a positive effect of iron supplementation in this cohort.

SCFAs propanoic and acetic acids are associated with geography specific fermentation: Relatively higher levels of these compounds were observed at 180m post lactulose ingestion indicating colonic fermentation⁸. These findings are supported by previous evidence showing SCFAs are the main metabolites produced in the colon by bacterial fermentation⁹.

Hydrogen sulphide (H₂S) was significantly decreased after 28 days of iron supplementation following lactulose

ingestion. H₂S is considered to be detrimental to gut health thus decreases in this compound is beneficial¹⁰. Please see talk 'The past, present and future of breath testing for bacterial overgrowth' at BBCon 2022 to hear more about the effects of H₂S and its role in this study.

It should also be noted that a limitation of this study was that blank measurements were done at the clinic site by drawing ambient air into a bag via a syringe. This air may have atmospheric contamination due to cleaning agents, perfumes etc. Thus, the room ambient and breath samples may be noisy. The statistically significant changes were calculated as two standard deviations from lab ambient.

Another limitation of this study was that not all subjects were healthy. After filling out clinical questionnaires, it was determined that subjects showed signs of small intestinal bacterial overgrowth (SIBO) or irritable bowel syndrome (IBS). These underlying conditions would have likely impacted VOCs produced.

References

- Mathers, C., Fat, D.M., Boerma, J.T. and World Health Organization eds., 2008. *The global burden of disease: 2004 update*. Geneva, Switzerland: World Health Organization.
- Schrier, S.L. and A., 2018. *Treatment of Iron Deficiency Anaemia in Adults*. [online]. Available at: <https://www.uptodate.com/contents/treatment-of-iron-deficiency-anemia-in-adults> [Accessed 21 Nov 2018].
- Camaschella, C., 2019. Iron deficiency. *Blood*, 133(1), pp.30-39.
- Goddard, A.F., James, M.W., McIntyre, A.S., Scott, B.B. and on behalf of the British Society of Gastroenterology, 2011. Guidelines for the management of iron deficiency anaemia. *Gut*, 60(10), pp.1309-1316.
- Rezaie A et al. Pimentel M. Hydrogen and Methane-Based Breath Testing in Gastrointestinal Disorders: The North American Consensus. *Am J Gastroenterol*. 2017 May;112(5):775-784. doi: 10.1038/sj.ajg.2017.46. Epub 2017 Mar 21. PMID: 28323273; PMCID: PMC5418558.
- Martin-Gallausiaux C, et al. SCFAs: mechanisms and functional importance in the gut. *Proc Nutr Soc*. 2021 Feb;80(1):37-49. doi: 10.1017/S002966520006916. Epub 2020 Apr 2. PMID: 32238208.
- Blaak EE, et al. Short chain fatty acids in human gut and metabolic health. *Brief Microbes*. 2020 Sep 1;1(5):411-455. doi: 10.3929/ethz-b-00057. Epub 2020 Aug 31. PMID: 32865024.
- Ghoshal UC. How to interpret hydrogen breath tests. *J Neurogastroenterol Motil*. 2011 Jul;17(3):312-7. doi: 10.5056/jnm.2011.17.3.312. Epub 2011 Jul 14. PMID: 21860825; PMCID: PMC3155069.
- Mortensen PB, Clausen MR. Short-chain fatty acids in the human colon: relation to gastrointestinal health and disease. *Scand J Gastroenterol Suppl*. 1996;236:132-48. doi: 10.3109/0036552960904568. PMID: 8726286
- Dordevic D et al. Hydrogen sulfide toxicity in the gut environment: Meta-analysis of sulfate-reducing and lactic acid bacteria in inflammatory processes. *Journal of Advanced Research*. Volume 27, 2021, Pages 55-69, ISSN 2090-1232, https://doi.org/10.1016/j.jare.2020.03.003.